

Accelerating cosmologies from non-local higher-derivative gravity

Salvatore Capozziello¹, Emilio Elizalde², Shin'ichi Nojiri³, Sergei D. Odintsov⁴

¹*Dipartimento di Scienze fisiche, Università di Napoli "Federico II", INFN Sez. di Napoli, Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, I-80126, Napoli, Italy*

²*Consejo Superior de Investigaciones Científicas ICE/CSIC-IEEC, Campus UAB, Facultat de Ciències, Torre C5-Parrell-2a pl, E-08193 Bellaterra (Barcelona) Spain*

³*Department of Physics, Nagoya University, Nagoya 464-8602, Japan*

⁴*Institució Catalana de Recerca i Estudis Avançats (ICREA) and Institut de Ciències de l'Espai (IEEC-CSIC), Campus UAB, Facultat de Ciències, Torre C5-Par-2a pl, E-08193 Bellaterra (Barcelona), Spain.*

We study accelerating cosmological solutions of a general class of non-linear gravities which depend on Gauss-Bonnet and other higher derivative invariants. To achieve this goal a local formulation with auxiliary scalars for arbitrary higher-derivative non-local gravity is developed. It is demonstrated that non-local Gauss-Bonnet gravity can be reduced, in the local formulation, to a model of string-inspired scalar-Gauss-Bonnet gravity. A natural unification, in the theory here developed, of the early-time inflation epoch with a late-time acceleration stage can also be realized.

PACS numbers: 04.50.+h, 04.80.Cc, 98.80.-k, 11.25.-w, 95.36.+x

1. Introduction. It has been seen recently that not only early-time inflation but also late-time acceleration may be naturally constructed in modified gravity (for a review of modified gravities, see [1], and for review of observational predictions from such theories, see [2]). It is remarkable that, moreover, a consistent unification of early-time inflation with the late-time dark energy universe can indeed be achieved in modified gravity (for a first such example, see [3]). However, the models under consideration represent local theories. At the same time, it is also well-known that quantum gravity and stringy considerations lead to a low-energy effective action which contains non-local, as well as local terms.

The proposal to use specific non-local gravity depending only on the scalar curvature as a dark energy source has been made in Refs. [4, 5]. Moreover, it has been demonstrated that different cosmological solutions including phantom and/or quintessence acceleration can be found, for specific versions of such non-local gravities [5, 6, 7]. In addition, the unification of early-time inflation with late-time acceleration in such theory, which turns out to be consistent with local tests, is also possible [5]. (A different approach to string-inspired non-local cosmology has been discussed in Ref. [8]). However, the structure of such non-local gravity in the local formulation with auxiliary fields is very complicated. In particular, an explicit de Sitter solution has been only found for a very restricted (exponential) class of models.

In the present paper we consider a more general class of non-local gravities which depend on Gauss-Bonnet and other higher-derivative invariants. First, we formulate the local version of the theory with auxiliary fields and show that almost all models of this sort admit the de Sitter cosmology solution. It is quite interesting to observe that a relation with string-inspired scalar-Gauss-Bonnet gravity (with possible account of higher-order string corrections) is found. The emergence of inflationary or late-time acceleration as well as the possibility to unify early-time inflation with late-time acceleration will be here established.

2. The de Sitter universe from non-local $\mathcal{G}$ -gravity. We now consider the following non-local model

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{\kappa^2}{2\alpha} \mathcal{G} \square^{-1} \mathcal{G} + \mathcal{L}_m \right). \quad (1)$$

Here $\mathcal{L}_m$ is the matter Lagrangian and $\mathcal{G}$ the Gauss-Bonnet invariant, namely $\mathcal{G} = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$. By introducing the scalar field ϕ , one may rewrite the action (1) in a local form:

$$S = \int d^4x \left(\frac{R}{2\kappa^2} - \frac{\alpha}{2\kappa^2} \partial_\mu \phi \partial^\mu \phi + \phi \mathcal{G} + \mathcal{L}_m \right). \quad (2)$$

In fact, from the ϕ -equation, we find $\phi = -\frac{\kappa^2}{\alpha} \square^{-1} \mathcal{G}$. By substituting this expression into (2), we reobtain (1). It is remarkable that the above action belongs to a specific class of string-inspired, scalar-Gauss-Bonnet gravity which is known to be consistent with the local tests and may serve as a realistic candidate for dark energy[9]. Unlike what happens in standard stringy compactifications, the scalar-GB coupling is not an exponential scalar function. The scalar potential is zero, as it should be in the low-energy, perturbative approach to the string effective action.

Let us now study the de Sitter solutions which may describe late-time universe acceleration in such non-local gravity. By assuming that the FRW metric with flat spatial part and ϕ only depend on the cosmological time, the equations

of motion are

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\alpha}{2\kappa^2}\dot{\phi}^2 - 24\dot{\phi}H^3 + \rho_m, \quad 0 = -\frac{\alpha}{\kappa^2}(\ddot{\phi} + 3H\dot{\phi}) + 24(\dot{H}H^2 + H^4). \quad (3)$$

Here ρ_m is the energy density of matter and H is the Hubble rate $H \equiv \dot{a}/a$. When the matter contribution is neglected, assuming $\dot{\phi}$ and H are constant $\dot{\phi} = c_\phi$, $H = H_0$, one can solve Eqs. (3) as follows: $c_\phi = \frac{8\kappa^2}{\alpha}H_0^3$, $H_0^4 = -\frac{3\alpha}{160\kappa^4}$. Therefore, if α is allowed to be negative, there is a de Sitter universe solution, namely

$$H = H_0 = \left(-\frac{3\alpha}{160\kappa^4}\right)^{1/4}. \quad (4)$$

On the other hand, a negative α means that the scalar field has a ‘wrong’ kinetic term, like a ghost or phantom. As we will see later in (13), the de Sitter space solution corresponds to the case $a \rightarrow \infty$ (late-time acceleration) when matter is included. Even at the early universe, before the matter was created, we can use this solution for the description of the inflationary epoch.

Note that the second equation (3) yields

$$\left(-\frac{\alpha}{\kappa^2}\dot{\phi} + 8H^2\right)a^3 = C. \quad (5)$$

Here C is a constant. We should note that the action (2) is invariant under a constant shift of the scalar field ϕ : $\phi \rightarrow \phi + c$ (c is a constant) since $\mathcal{G}$ is a total derivative. The quantity C is the Noether charge corresponding to the invariance. By using (5), we may delete $\dot{\phi}$ in the first equation in (3) and obtain,

$$0 = -\frac{3H^2}{\kappa^2} - \frac{160\kappa^2H^6}{\alpha} + \frac{16\kappa^2CH^3}{\alpha a^3} + \frac{\kappa^2C^2}{\alpha a^6} + \rho_m. \quad (6)$$

When ρ_m is given by the sum of the contributions from all forms of matter with constant EoS parameter w_i , one gets $\rho_m(a) = \sum_i \rho_{0i}a^{-3(w_i+1)}$. Then we can solve algebraically Eq. (6) with respect to H as a function of a : $H = \mathcal{H}(a)$. By the definition of H , the equation has the form as $\dot{a} = a\mathcal{H}(a)$. Then we find $t = \int \frac{da}{a\mathcal{H}(a)}$, which gives, at least formally, the solution of (6) and therefore the FRW universe.

As a simple case, we may consider the case $C = 0$. Then Eq. (6) reduces to the following form

$$0 = \mathcal{F}(x) \equiv x^3 + \frac{3\alpha}{160\kappa^4}x - \frac{\alpha\rho_m(a)}{160\kappa^2}. \quad (7)$$

Here $x \equiv H^2$ and therefore $x \geq 0$. Since $\mathcal{F}'(x) = 3\left(x^2 + \frac{\alpha}{160\kappa^4}\right)$, when $\alpha > 0$, $\mathcal{F}(x)$ is a monotonically increasing function of x . On the other hand, when $\alpha < 0$, we find

$$\mathcal{F}'(x_\pm) = 0, \quad \mathcal{F}(x_\pm) = \mp \left(-\frac{\alpha}{160\kappa^4}\right)^{3/2} - \frac{\alpha\rho_m}{160\kappa^2}, \quad x_\pm \equiv \pm \left(-\frac{\alpha}{160\kappa^4}\right)^{1/2}. \quad (8)$$

Since

$$\mathcal{F}(0) = -\frac{\alpha\rho_m}{160\kappa^2}, \quad (9)$$

when $\alpha > 0$, since $\mathcal{F}(0) < 0$ and $\mathcal{F}(x)$ is a monotonically increasing function, there is only one positive solution in (7):

$$H^2 = x = \alpha_+ + \alpha_- , \quad \alpha_\pm^3 = \frac{\alpha\rho_m}{320\kappa^2} \left(1 \pm \sqrt{1 + \frac{\alpha}{40\kappa^8\rho_m^2}}\right). \quad (10)$$

We now consider the case that ρ_m comes from a unique kind of matter, with constant EoS parameter w . Then, when $a \rightarrow 0$, one obtains $\alpha_+^3 \rightarrow \frac{\alpha\rho_0}{160\kappa^2}a^{-3(1+w)}$ and $\alpha_- \rightarrow 0$. Therefore, $H \propto a^{-(1+w)/2}$ and we find $a \propto t^{2/(1+w)}$, which could be compared with the usual FRW universe in Einsteinian gravity, where $a \propto t^{2/3(1+w)}$. On the other hand, when $a \rightarrow \infty$, Eq. (10) implies that $H \propto a^{-3(1+w)/2}$ and therefore $a \propto t^{2/3(1+w)}$, which reproduces the result in Einstein’s gravity. This indicates that such non-local term becomes less dominant as compared with the Einstein-Hilbert term at the late universe epoch.

We may consider the case when $\alpha < 0$. Since $\mathcal{F}(0) > 0$ in this case, if $\mathcal{F}(x_+) > 0$, there is no positive solution for x . On the other hand, if $\mathcal{F}(x_+) < 0$, that is, when

$$-\alpha > 40\kappa^6\rho_m^2, \quad (11)$$

there are two positive solutions and one negative solution in (7), namely

$$x = \tilde{\alpha}_+ + \tilde{\alpha}_-, \quad \tilde{\alpha}_+\xi + \tilde{\alpha}_-\xi^2, \quad \tilde{\alpha}_+\xi^2 + \tilde{\alpha}_-\xi, \quad \tilde{\alpha}_\pm^3 = \frac{\alpha\rho_m}{320\kappa^2} \left(1 \pm i\sqrt{-1 - \frac{\alpha}{40\kappa^8\rho_m^2}} \right). \quad (12)$$

Eq. (11) indicates that there is a lower bound in a as $a \geq a_{\min} \equiv \left(-\frac{\alpha}{40\kappa^4\rho_0^2} \right)^{1/6(1+w)}$. Here it is assumed that $w > -1$. We should note that even if $a = a_{\min}$, x and therefore H do not vanish, which means that the existence of $a_{\min}$ does not imply the bounce solution. It is likely that a universe with $a = a_{\min}$ could appear, say, as a quantum correction.

When a is large, we find that $\mathcal{F}(0)$ in (9) vanishes and $\alpha_\pm$ tends to a constant: $\alpha_\pm^3 \rightarrow \pm i \frac{1}{640\kappa^6} \sqrt{-\frac{\alpha^3}{10}}$, which gives

$$H^2 = x = 0, \quad \pm \sqrt{-\frac{3\alpha}{160\kappa^4}}. \quad (13)$$

The positive solution of x in (13) corresponds to the de Sitter solution (4).

When the Einstein-Hilbert term in (1) or (2) can be neglected, by assuming

$$\phi = \frac{a_\phi}{t^2}, \quad H = \frac{h_0}{t} \quad (a = a_0 t^{h_0}), \quad (14)$$

Eqs. (3) reduce to the following algebraic equations

$$0 = a_\phi \left(\frac{\alpha}{\kappa^2} a_\phi + 24h_0^3 \right), \quad 0 = (h_0 - 1) \left(\frac{\alpha}{\kappa^2} a_\phi + 4h_0^3 \right). \quad (15)$$

The equations (15) have a non-trivial solution ($a_\phi \neq 0$ and $h_0 \neq 0$): $h_0 = 1$ and $a_\phi = -\frac{24\kappa^2}{\alpha}$.

We can now include matter. Then, the first equation in (15) is modified as

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\alpha}{2\kappa^2} \dot{\phi}^2 - 24\dot{\phi}H^3 + \rho_m. \quad (16)$$

When one may neglect the Einstein-Hilbert term in (1) or (2) again, and if we assume (14), there is a non-trivial solution when $h_0 = \frac{4}{3(w+1)}$. In fact, Eqs. (3) reduce to the following algebraic expressions

$$0 = \frac{2\alpha}{\kappa^2} a_\phi^2 + \rho_0 a_0^{-3(w+1)} + 48a_\phi \left(\frac{4}{3(1+w)} \right)^3, \quad 0 = \left(\frac{4}{3(1+w)} - 1 \right) \left(\frac{\alpha}{\kappa^2} a_\phi + 4 \left(\frac{4}{3(1+w)} \right)^3 \right). \quad (17)$$

If the matter is radiation with $w = 1/3$, the second equation is satisfied and the first equation acquires the following form: $0 = \frac{2\alpha}{\kappa^2} a_\phi^2 + \rho_0 a_0^{-4} + 48a_\phi$. Then, if $\tilde{D} \equiv (24)^2 - \frac{2\alpha\rho_0}{\kappa^2 a_0^4} > 0$, there are non-trivial solutions: $a_\phi = \frac{\kappa^2}{2\alpha} \left(-24 \pm \sqrt{\tilde{D}} \right)$.

On the other hand, when $w \neq 1/3$, we find $a_\phi = -\frac{4\kappa^2}{\alpha} \left(\frac{4}{3(1+w)} \right)^3$, $\rho_0 a_0^{-4} = \frac{160\kappa^2}{\alpha} \left(\frac{4}{3(1+w)} \right)^6$. The second equation requires that α should be positive. Thus, a phantom-like universe solution from purely non-local gravity, without the R -term, has been found.

We now consider another version of (1), that is,

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{\kappa^2}{2\alpha} \left(\frac{1}{\mathcal{G}} \right) \square^{-1} \left(\frac{1}{\mathcal{G}} \right) + \mathcal{L}_m \right). \quad (18)$$

Here α is a dimensional constant. Naively this action could be relevant in the late universe stage, where the curvature, and therefore $\mathcal{G}$, is small ($1/\mathcal{G}$ is large) owing to the non-locality of the action (18). However, there is a possibility that the action could be relevant even for the early universe. By introducing three scalar fields, ϕ , ξ , and η , one may rewrite the action (18) under the following form:

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} + \frac{\alpha}{2} \partial_\mu \phi \partial^\mu \phi + \frac{\alpha}{\kappa^7} \frac{\phi}{\eta} + \xi(\eta - \mathcal{G}) \right). \quad (19)$$

Again, the action reminds of string-inspired gravity, including the first Gauss-Bonnet correction. However, it may correspond to some complicated compactification and/or to the contribution of non-perturbative terms, because of the absence of kinetic terms for the two scalars and the appearance of a non-trivial multi-scalar potential.

In the FRW universe, the action leads to the following equations

$$0 = -\frac{3}{\kappa^2}H^2 - \frac{\alpha}{2}\dot{\phi}^2 + 24\dot{\xi}H^3, \quad 0 = \alpha\left(\ddot{\phi} + 3H\dot{\phi}\right) + \frac{\alpha}{\kappa^7\eta}, \quad 0 = -\frac{\alpha}{\kappa^2}\frac{\phi}{\eta^2} + \xi, \quad 0 = \eta - 24\left(H^4 + \dot{H}H^2\right). \quad (20)$$

By assuming $\phi = c_\phi t$, $\eta = \eta_0$, $\xi = \xi_0 t$, and $H = H_0$, with constants c_ϕ , η_0 , ξ_0 , and H_0 , the equations in (20) reduce to the algebraic ones

$$0 = -\frac{3}{\kappa^2}H_0^2 - \frac{\alpha}{2}c_\phi^2 + 24\xi_0H_0^3, \quad 0 = 3\alpha H_0 c_\phi + \frac{\alpha}{\kappa^7\eta_0}, \quad 0 = -\frac{\alpha}{\kappa^2}\frac{c_\phi}{\eta_0^2} + \xi_0, \quad 0 = \eta_0 - 24H_0^4. \quad (21)$$

We can solve Eqs. (21) with respect to c_ϕ , ξ_0 , and η_0 as follows: $c_\phi = -\frac{1}{72\kappa^7 H_0^3}$, $\xi_0 = -\frac{\alpha}{3(24)^3 \kappa^{14} H_0^{13}}$, $\eta_0 = 24H_0^4$, and $H_0^{12} = -\frac{7}{54(24)^2 \kappa^{12}}$. Therefore if α is negative, there could be a de Sitter universe solution, where H is a constant: $H_0 = \left(-\frac{7}{54(24)^2}\right)^{1/12} \frac{1}{\kappa}$. As in the case of (4), the de Sitter solution may both describe late time acceleration or inflation in the early universe. Hence, we have demonstrated that non-local G -gravities of different types may lead to an alternative description of dark energy and/or inflationary dynamics. Of course, for realistic description some version of the theory should be elaborated in more detail.

3. General non-linear higher-derivative gravity. We can now consider a generalization of (1) which depends on all three curvature tensors (the case of two different functions F may be considered by analogy)

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{1}{2\beta} F(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \square^{-1} F(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right). \quad (22)$$

Here $F(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho})$ is a scalar function of the scalar curvature R , the Ricci tensor $R_{\mu\nu}$, and the Riemann tensor. Introducing an auxiliary field ϕ as in (2), the action (22) can be rewritten in a local form

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{\beta}{2} \partial_\mu \phi \partial^\mu \phi - \phi F(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right). \quad (23)$$

Even if the action includes higher powers of the inverse of $\square$ (compare with [7]) as

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \sum_{n=1}^N \frac{1}{2\beta_n} F_n(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \square^{-n} F_n(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right), \quad (24)$$

we can still rewrite the action (24) in a local form by introducing auxiliary scalar fields (compare with the case of on-local R -gravity [7]).

One may consider a more complicated generalization, as

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \square^{-1} F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right). \quad (25)$$

Here we assume that F_1 is a function different from F_2 . By properly multiplying the Planck scale M_{Pl} , we may set the mass dimensions of F_1 and F_2 to be 3. Then, the above action can be written in the following form:

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{1}{2} (F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) + F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho})) \square^{-1} (F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) + F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho})) \right. \\ \left. + \frac{1}{2} F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \square^{-1} F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) + \frac{1}{2} F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \square^{-1} F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right). \quad (26)$$

By introducing now three scalar fields, ϕ_1 , ϕ_2 , ϕ_3 , we can rewrite (26) in a local form as

$$S = \int d^4x \left(\frac{R}{2\kappa^2} + \frac{1}{2} \partial_\mu \phi_1 \partial^\mu \phi_1 + \frac{1}{2} \partial_\mu \phi_2 \partial^\mu \phi_2 \right. \\ \left. - \frac{1}{2} \partial_\mu \phi_3 \partial^\mu \phi_3 - (\phi_1 + \phi_3) F_1(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) - (\phi_2 + \phi_3) F_2(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) \right). \quad (27)$$

Note that if F_1, F_2 include Gauss-Bonnet and higher-order terms which are cubic and fourth-order in the curvature invariants, the above theory represents the low-energy string effective action with higher-order corrections. The de Sitter solutions for such effective action have been studied in [10].

As an example of (22) or (23), we may consider the case $F(R, R_{\mu\nu}, R_{\mu\nu\sigma\rho}) = c_1\mathcal{G} + c_2R$, with constants c_1 and c_2 . Then, in the FRW metric, we find the following equations:

$$0 = -\beta \left(\ddot{\phi} + 3H\dot{\phi} \right) - 24c_1 \left(\dot{H}H^2 + H^4 \right) - 6c_2 \left(\dot{H} + 2H^2 \right) , \quad (28)$$

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\beta}{2} + 24c_1\dot{\phi}H^3 - 3c_2 \left(\phi H^2 - \dot{\phi}H \right) + \rho_m . \quad (29)$$

When the contribution from matter can be neglected, that is, provided $\rho_m = 0$, if c_2/c_1 is negative there is a de Sitter space solution: $H = \sqrt{-\frac{c_2}{2c_1}}$ and $\phi = -\frac{1}{c_2\kappa^2}$. As we have neglected the contribution from matter, again, the de Sitter solution could be valid to describe late time acceleration or inflation before matter was created in the very early universe.

Now, the following model can be considered:

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} + (c_1\mathcal{G} + c_2R) f(\Box^{-1}(c_1\mathcal{G} + c_2R)) \right\} . \quad (30)$$

Here c_1 and c_2 are constant. By introducing two scalar fields, ξ and ϕ , we may rewrite the action (30) in a local form

$$S = \int d^4x \sqrt{-g} \left\{ \frac{R}{2\kappa^2} - (\xi - f(\phi))(c_1\mathcal{G} + c_2R) - \partial_\mu \xi \partial^\mu \phi \right\} . \quad (31)$$

One may check that the action (31) is equivalent to the action (30). By variation of ξ and ϕ , we have

$$0 = \Box \phi - (c_1\mathcal{G} + c_2R) , \quad (32)$$

$$0 = \Box \xi + f'(\phi)(c_1\mathcal{G} + c_2R) . \quad (33)$$

If we delete ϕ in (31) by using (32), we reobtain the action (30). In the FRW background, Eqs. (32) and (33) can be rewritten as

$$0 = -\left(\ddot{\phi} + 3H\dot{\phi} \right) - \left\{ 24c_1 \left(\dot{H}H^2 + H^4 \right) + 6c_2 \left(\dot{H} + 2H^2 \right) \right\} , \quad (34)$$

$$0 = -\left(\ddot{\xi} + 3H\dot{\xi} \right) + f'(\phi) \left\{ 24c_1 \left(\dot{H}H^2 + H^4 \right) + 6c_2 \left(\dot{H} + 2H^2 \right) \right\} . \quad (35)$$

On the other hand, the equation corresponding to the FRW one has the following form

$$0 = -\frac{3}{\kappa^2}H^2 + 24c_1 \left(\dot{\xi} - f'(\phi)\dot{\phi} \right) H^3 + 3c_2 \left\{ (\xi - f(\phi))H^2 - \left(\dot{\xi} - f'(\phi)\dot{\phi} \right) H \right\} + \rho_m . \quad (36)$$

We now investigate if there could be a de Sitter space solution. We assume H is a constant $H = H_0$ and neglect matter, $\rho_m = 0$. Then, Eq. (34) can be integrated to be

$$\phi = -\left(8c_1H_0^3 + 4c_2H_0 \right) t - \phi_1 e^{-3H_0 t} + \phi_2 . \quad (37)$$

Here ϕ_1 and ϕ_2 are constants of integration. For simplicity, we consider the case $\phi_1 = \phi_2 = 0$ and we assume that $f(\phi)$ has the following form:

$$f(\phi) = f_0 e^{b\phi} . \quad (38)$$

Here b is a constant. By substituting (37) with $\phi_1 = \phi_2 = 0$ into (38) and integrating (35), we obtain

$$\xi = -\frac{3H_0 f_0}{3H_0 - \beta} e^{-\beta t} - \frac{\xi_1}{3H_0} e^{-3H_0 t} + \xi_2 . \quad (39)$$

Here ξ_1 and ξ_2 are constants of integration and β is defined by

$$\beta \equiv \left(8c_1H_0^3 + 4c_2H_0 \right) b . \quad (40)$$

Substituting now (37) with $\phi_1 = \phi_2 = 0$ and (40) into (36), one gets

$$0 = -3H_0^2 \left(\frac{1}{\kappa^2} + c_2 \xi_2 \right) + \frac{(24c_1H_0^3 + 3c_2H_0)\beta^2 - 3c_2H_0^2\beta}{3H_0 - \beta} f_0 e^{-\beta t} + (24c_1H_0^3 - 4c_2H_0) \xi_1 e^{-3H_0 t} . \quad (41)$$

When $\beta \neq 3H_0$ nor $\beta \neq 0$, Eq. (41) yields

$$\xi_2 = -\frac{1}{c_2 \kappa^2} , \quad (42)$$

and

$$0 = 6c_1 H_0^2 - c_2 \quad \text{or} \quad \xi_1 = 0 . \quad (43)$$

Furthermore,

$$\beta = -1 - \frac{8c_1 H_0}{c_2} \quad \text{or} \quad b = -\frac{1 + \frac{8c_1 H_0}{c_2}}{8c_1 + 4c_2 H_0} . \quad (44)$$

Then if we choose b in (38) as in (44), there is a de Sitter solution with $H = H_0$. When $c_1 = 0$, that is, without the Gauss-Bonnet term, we have $\xi_1 = 0$ and (42). Eq. (41) also gives $\beta = -1$, which corresponds to the solution in [5]. In the same way as in Ref. [5] one can suggest the unification of early-time inflation with late-time acceleration within the theory. It is also remarkable that unlike in the case of non-local gravities with R -dependence only, the above theory is very rich and does admit de Sitter universe solutions for a variety of actions.

The unification of late-time acceleration and inflation in the early universe can in fact be easily achieved by adding a $F(R)$ -term to the action of non-local Gauss-Bonnet gravity (1), as

$$S = \int d^4x \sqrt{-g} \left(\frac{R}{2\kappa^2} - \frac{\kappa^2}{2\alpha} \mathcal{G} \square^{-1} \mathcal{G} + F(R) + \mathcal{L}_m \right) , \quad (45)$$

or to the action

$$S = \int d^4x \left(\frac{R}{2\kappa^2} - \frac{\alpha}{2\kappa^2} \partial_\mu \phi \partial^\mu \phi + \phi \mathcal{G} + F(R) + \mathcal{L}_m \right) . \quad (46)$$

Here $F(R)$ is a proper function of the scalar curvature R (the above theory has been proposed for dark energy description in Ref. [11]). Then the first equation in Eq. (3) is changed as

$$0 = -\frac{3H^2}{\kappa^2} + \frac{\alpha}{2\kappa^2} \dot{\phi}^2 - 24\dot{\phi}H^3 - F(R) + 6(H^2 + \dot{H})F'(R) - 36(4H^2\dot{H} + H\ddot{H})F''(R) + \rho_m , \quad (47)$$

but the second equation is not changed and therefore we have (5) again.

One can consider the case where the early inflation could be mainly generated by the $F(R)$ -term and the late-time accelerating expansion could be generated by the non-local Gauss-Bonnet term. We now choose $F(R) = \beta R^2$. Here β is a constant. We choose α to be very small and ϕ starts with $\phi = 0$. Hence, at the early universe the non-local Gauss-Bonnet term is very small and could be neglected. We also assume that the contribution from matter could be also neglected in the early universe. Then due to the $F(R)$ -term, there occurs R^2 -inflation. After the end of inflation, the radiation/matter dominance era takes over, where H behaves as $H \sim h_0/(t+t_0)$. Then by integrating (5), we find

$$\phi = \phi_0 + \frac{\kappa^2}{\alpha} \left(-\frac{8h_0^2}{t+t_0} - \tilde{C}(t+t_0)^{-3h_0+1} \right) . \quad (48)$$

Here ϕ_0 and $\tilde{C}$ are constants of integration, which could be determined by proper initial conditions. When t becomes large, the value of ϕ goes to ϕ_0 . If ϕ_0 can be large enough, the non-local Gauss-Bonnet term can be dominated and the universe goes to asymptotically de Sitter space, which describes late-time acceleration. Thus, a mixture of non-local Gauss-Bonnet gravity with local modified gravity, on top of being a rather natural theory made up of the fundamental ingredients one has at hand, yields a very realistic model for the unification of early-time inflation with late-time acceleration.

4. Discussion.

In summary, we have demonstrated in this paper that a large class of non-local higher-derivative gravities admits accelerating cosmologies which can naturally describe inflation, dark energy and/or the wholly unified inflation-radiation/matter dominance-dark energy, in the universe expansion history. We have chiefly concentrated on models which depend on a Gauss-Bonnet invariant but this was essentially for technical reasons: the simplification of the equations of motion. There is no problem to extend the study to account for Riemann squared tensor- and/or Ricci squared tensor-dependent non-local gravity. In addition, non-linear GB gravity can be represented as a specific

string-inspired gravity, what hints towards a fundamental origin from string theory. Unlike in the case of pure curvature-dependent non-linear gravity, many different versions of the theory under considerations exhibit de Sitter universe solutions which maybe used to describe the different accelerating epochs. It is known that such solutions represent the fixed points of the cosmological dynamics where the effective equation of state parameter is $w = -1$. Moreover, the stability of the fixed points can be studied analytically or numerically. Hence, it is straightforward to construct the effective phantom and quintessence-like cosmologies which lie below or above the corresponding de Sitter fixed points in a similar way as in Ref. [7].

In order to select the most realistic version from the above theories, local tests, PPN analysis and accurate observational data must be used. For instance, cosmologies derived from non-local higher-derivative gravity should be experimentally tested in order to make a short list of most viable theories. From a genuine cosmological viewpoint, several non-local models give rise to very similar dynamics. A way to overcome such a shortcoming could be to search for signatures in the stochastic background of gravitational waves in order to discriminate and constrain models. In fact, the production of primordial gravitational waves, if revealed, is a robust prediction for any model attempting to describe the cosmological evolution at early epochs. It may be directly derived from the inflationary scenario [13], which fits the WMAP data well, being the agreement particularly good with exponential inflation and the spectral index $\simeq 1$. The main characteristics of the gravitational background produced by cosmological sources depend both on the emission properties of each single source and on the source rate evolution with the redshift. To this purpose, one can take into account the primordial physical process which gave rise to a characteristic spectrum Ω_{sgw} for the early stochastic background of relic scalar gravitational waves by which we can recast the further degrees of freedom coming from non-local higher derivative gravity. This approach can greatly contribute to constrain the viable cosmological models. The physical process related to the production has been analyzed for the first several tensorial components of standard General Relativity. Actually the process can be improved considering also the further scalar-tensor components strictly related to the degrees of freedom of non-local gravity. It should be emphasized that the stochastic background of scalar gravitational waves can be described in terms of scalar fields ϕ and characterized by a dimensionless spectrum. In our case, considering a generic non-local theory of order $(2n + 2)$ as, for example,

$$\mathcal{L} = \sqrt{-g}F(R, \square R, \dots \square^k R, \square^n R), \quad (49)$$

it is straightforward to show that, by linearizing the theory, one obtains

$$\prod_{k=0}^n (\square - m_k^2) \phi_k = 0, \quad (50)$$

with n massive scalar fields (m_k real) or tachyons (m_k imaginary) conformally related to the curvature invariants. The interferometric space experiment LISA could put tight constraints on such masses. An analysis similar to the one in [12] will be developed for the theory under consideration elsewhere.

Acknowledgments This research was supported by a INFN-CICYT bilateral project, by *Azione Integrata Italia-Spagna 2007* (MIUR Prot. No. 464, 13/3/2006) grant, and by MCIN (Spain) projects FIS2006-02842 and PIE2007-501023. The work by S.N. is supported by Min. of Education, Science, Sports and Culture of Japan under grant no. 18549001 and Global COE Program of Nagoya University provided by the Japan Society for the Promotion of Science (G07).

-
- [1] S. Nojiri and S. D. Odintsov, arXiv:hep-th/0601213.
 - [2] S. Capozziello and M. Francaviglia, Gen.Rel.Grav. **40**, 357 (2008), arXiv:0706.1146[astro-ph].
 - [3] S. Nojiri and S. D. Odintsov, Phys. Rev. D **68**, 123512 (2003) [arXiv:hep-th/0307288].
 - [4] S. Deser and R. Woodard, Phys. Rev. Lett. **99**, 111301 (2007).
 - [5] S. Nojiri and S. D. Odintsov, Phys. Lett. B **659**, 821 (2008) [arXiv:0708.0924 [hep-th]].
 - [6] T. Koivisto, Phys. Rev. D **77**, 123513 (2008); [arXiv:0807.3778[gr-qc]].
 - [7] S. Jhingan, S. Nojiri, S. D. Odintsov, M. Sami, I. Thongkool and S. Zerbini, Phys. Lett. B **663**, 424 (2008) [arXiv:0803.2613 [hep-th]].
 - [8] I. Arefeva, L. Joukovskaya and S. Vernov, JHEP **0707**, 087 (2007).
 - [9] S. Nojiri, S. D. Odintsov and M. Sasaki, Phys. Rev. D **71**, 123509 (2005) [arXiv:hep-th/0504052].
 - [10] E. Elizalde, S. Jhingan, S. Nojiri, S. D. Odintsov, M. Sami and I. Thongkool, Eur. Phys. J. C **53**, 447 (2008) [arXiv:0705.1211[hep-th]]; K. Maeda and N. Ohta, Phys. Rev. D **71**, 063520 (2005); K. Akune, K. Maeda and N. Ohta, Phys. Rev. D **73**, 103506 (2006); M. Iihoshi and S. Ketov, arXiv:0707.3359[hep-th].
 - [11] S. Nojiri, S. D. Odintsov and P. Tretyakov, Phys. Lett. B **651**, 224 (2007) [arXiv:0704.2520[hep-th]].
 - [12] S. Capozziello, M. De Laurentis, S. Nojiri and S. D. Odintsov arXiv:0808.1335 [hep-th].
 - [13] G. S. Watson- "An exposition on inflationary cosmology" - North Carolina University Press (2000).